

****5/6/04 DRAFT****

**Fire Regime Condition Class (FRCC) Interagency Handbook
Reference Conditions**

Modelers: Wendel Hann, Reese **Date:** 5/6/04 **PNVG Code:** OCWI
Lolley, Cecilia McNicoll

Potential Natural Vegetation Group: Oak & Conifer Woodlands Interior
Southwest

Geographic Area: Southern Arizona, southern New Mexico, western Texas.

Description: Vegetation type is often referred to as being within the Madrean province; these landscapes are dominated by open evergreen oaks, alligator bark junipers, and Mexican pines ranging from 15 to 50 feet high, with a grass-dominated understory; type usually occupies foothills and mountains ranging from about 3000 to 7000 ft (1200-2200 m) elevation, and occurs in a zone between the warmer and dryer Plains Mesa Grassland at lower elevations and the moister Woodland-Grassland Complex on slopes/mesas at higher elevations; climate generally consists of mild winters and wet summers with mean annual precipitation ranging from about 10 to 25 inches; half of the precipitation typically occurs in summer, with the remainder occurring during winter/spring.

Fire Regime Description: Regime I (frequent surface- and mixed severity fires) with fire intervals generally ranging from 5-20 years long (approx. 10-yr MFI; Brown and Smith 2000, USDA 2002). Large-diameter alligator junipers and evergreen oaks often survive 1 to 3 low intensity fires resulting in “cat face” scars with char at the base of the tree, whereas Mexican pines can survive multiple low intensity fires. Fire severity can be mixed in both space and time, for example, high-severity fires can occur on relatively productive sites, or during extreme fire weather and prolonged droughts.

Vegetation Type and Structure

<i>Class</i>	<i>Percent of Landscape</i>	<i>Description</i>
A: Early-seral post-fire grass-forb	10	Post-fire grass and fire-adapted forbs: herbaceous life form with 10-30 % canopy and 20% average; mountain muly, blue grama, sideoats grama, asters, penstemons, sprouting shrubs
B: Mid-seral tree-shrub/grass	20	Mid-seral woodland, typically in more productive draws and northerly aspects: woodland life form with 15-70% canopy, average of 55%; alligator juniper, oaks, mahogany, mountain muly, blue grama
C: Mid-seral open grass with scattered shrubs and trees	25	Mid-seral grasslands on southerly slopes & ridges: grass dominated herbaceous life form with species such as mountain muhly, blue grama, and sideoats grama; 25-65% herbaceous cover; 5-15% canopy of scattered

D: Late-seral open tree-shrub/grass	30	trees and shrubs, such as alligator juniper, oaks; Late-seral open woodland on slopes & ridges: woodland life form with 5-35% canopy, 25% average; alligator juniper, oaks, mountain mully, blue grama, sideoats grama;
E: Late-seral closed tree-shrub/grass	15	Late-seral closed woodland typically in draws or on steep rocky or thin soil slopes & ridges: woodland life form with 35-70% canopy, average of 55%;alligator juniper, oaks, mahogany, scattered shrubs and grasses
Total	100	

Fire Frequency and Severity

<i>Fire Frequency-Severity</i>	<i>Modeled Probability</i>	<i>Pct, All Fires</i>	<i>Description</i>
Replacement Fire	.03	33	Primarily stand replacmt. in E and B
Non-Replacement Fire	.062	67	Primarily maintenance in A, C, and D
All Fire Frequency*	.092	100	

*Sum of replacement fire and non-replacement fire probabilities.

References

- Bailey, Robert G. 1995. Descriptions of the ecoregions of the United States. 2nd ed. Rev. and expanded (1st ed. 1980). Misc. Publ. No. 1391 (rev.), Washington DC: USDA Forest Service. 108 p. with separate map at 1:7,500,000.
- Brown, David E. 1994. Biotic communities southwestern United States and northwestern Mexico. University of Utah Press, Salt Lake City, UT. 342 p.
- Brown, James K.; Smith, Jane Kapler, eds. 2000. Wildland fire in ecosystems: effects of fire on flora. Gen. Tech. Rep. RMRS-GTR-42-vol. 2. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 257 p.
- Kuchler, A. W. 1964. Manual to accompany the map of potential natural vegetation of the conterminous United States. American Geographical Society. Spec. Publ. No. 36. Lib. Congress Cat. Card Num. 64-15417. 156 p.
- Dick-Peddie, William A. 1993. New Mexico vegetation past present and future. University of New Mexico Press, Albuquerque, NM. 244 p.
- McNab, W. Henry; Avers, Peter E. 1994. Ecological subregions of the United States: section descriptions. USDA Forest Service, Ecosystem Management, Washington DC. WO-WSA-5. 250 p plus appendices and maps.
- Schmidt, Kirsten M, Menakis, James P., Hardy, Colin C., Hann, Wendel J., Bunnell, David L. 2002. Development of coarse-scale spatial data for wildland fire and fuel management. Gen. Tech. Rep. RMRS-GTR-87. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 41 p. + CD.
- U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (2002, December). Fire Effects Information System, [Online]. Available: <http://www.fs.fed.us/database/feis/>.